arXiv:q-alg/9610010v1 8 Oct 1996

q-alg/9610010
October, 1996

THE QUANTUM SYMPLECTIC CAYLEY-KLEIN GROUPS

N.A. GROMOV, I.V. KOSTYAKOV and V.V. KURATOV
Department of Mathematics, Komi Science Centre,
Ural Division, Russian Academy of Sciences,
Syktyvkar, 167000, Russia
E-mail : parma@omkomi.intec.ru

Abstract

The contraction method applied to the construction of the nonsemisimple quantum symplectic Cayley-Klein groups $Fun(Sp_q(n;j))$. This groups has been realised as Hopf algebra of the noncommutative functions over the algebra with nilpotent generators. The dual quantum algebras $sp_q(n;j)$ are constructed.

1 Introduction

A description of nonsemisimple quantum symplectic groups and algebras is suggested. It is known, that the groups motion of Cayley-Klein space, can be described uniformly [1], if the parameter $j, (j = 1, \iota$, where $\iota \neq 0, \iota^2 = 0)$ is introduced to the group. The symplectic groups is obtained by this method will be denoted as $Sp(n;j)$, and this introduction of parameter j shall be called the Cayley-Klein structure introduction on the group. The Cayley-Klein structure on the quantum symplectic group is defined and the description of nonsemisimple quantum symplectic groups is obtained. We used the notation [2].

2 Symplectic algebra sp(n;j)

Consider the space $\mathrm{R^n}(j) \otimes \mathrm{R^n}(j)$, which is obtained from $2n$-dimensional Euclidean space $\mathrm{R^n} \otimes \mathrm{R^n}$ by mapping

$$\psi : \ \mathrm{R^n} \otimes \mathrm{R^n} \rightarrow \mathrm{R^n}(j) \otimes \mathrm{R^n}(j),$$

$$\psi x_k = x_k J_k, \quad \psi x_{k'} = x_{k'} J_k, \quad k' = 2n+1-k, \ J_k = \prod_{m=1}^{k-1} j_m, \ k = 1, 2 \ldots n.$$

Group $Sp(n;j)$ is defined as the group of transformation of $2n-$dimensional space $R^n(j)\otimes R^n(j)$, preserving the bilinear form

$$(x,y)=\sum_{k=1}^{n}(x_k y_{k'}-x_{k'}y_k)J_k^2. \tag{1}$$

Cartesian coordinates $x_k, y_k,\ k=1,2\ldots n$, belongs to the first, and $x_{k'}, y_{k'}$— to the second factor in the direct product of spaces. For j-structure to be preserved, parameter j must be introduced in matrix of transformation $T(j)=(T_{ij})_{i,j=1}^{n}$ as follows

$$T_{ij}=J_m^k t_{ij},\ where\ \ k=i\ \ if\ \ i\le n;\quad k=i',\ \ i>n;\quad m=j,\ \ j\le n;$$

$$m=j',\ \ j>n,\qquad J_m^k=\prod_{n=min(k,m)}^{max(k,m)-1} j_n.$$

The matrix elements of T satisfies the additional relation

$$T^t C_0 T=C_0. \tag{2}$$

The elements of matrix C_0 are as follows $(C_0)_{ij}=\varepsilon_i\delta_{ij},\ \varepsilon_i=1$ if $i=1,2\ldots n$ and $\varepsilon_i=-1$ for $i=n-1,\ldots,2n$. The general element of the Shevalley basis of symplectic algebra $sp(n;j)$ appears as

$$H=\begin{pmatrix} h_1 & j_1x_1^+ & \cdot & \cdots & \cdots & \cdot \\ j_1x_1^- & h_2 & j_2x_2^+ & \cdots & \cdots & \cdot \\ \cdot & j_1x_2^+ & \cdots & \cdots & \cdots & \cdot \\ \cdot & \cdots & \cdots & \cdots & \cdots & \cdots \\ \cdot & \cdots & \cdots & -j_2x_2^- & -h_1 & -j_1x_1^+ \\ \cdot & \cdots & \cdots & \cdots & -j_1x_1^- & -h_2 \end{pmatrix}. \tag{3}$$

Choosing the generators of algebra $sp(n;j)$ in the following form

$$\begin{aligned} H_i &= e_{ii}-e_{i'i'}-e_{i+1,i+1}+e_{(i+1)',(i+1)'},\quad H_n=e_{nn}-e_{n',n'},\\ X_i^+ &= j_i(e_{i,i+1}-e_{(i+1)',i'}),\quad X_i^-=j_i(e_{i+1,i}-e_{(i)',(i+1)'}),\\ X_n^+ &= j_n e_{n,n+1},\quad X_n^-=j_n e_{n+1,n},\quad i=1,\ldots,n-1, \end{aligned} \tag{4}$$

where $(e_{ij})_{km}=\delta_{ik}\delta_{jm}$, the cmmutation relation for the Shevalley basis $sp(n;j)$ looks as

$$\left[H_i,X_j^{\pm}\right]=\pm A_{ij}X_j^{\pm},\quad \left[X_i^+,X_j^-\right]=\delta_{ij}j_i^2 H_j, \tag{5}$$

where A_{ij} is a Cartan matrix — $A_{ii}=2,\ A_{i,i-1}=A_{i-2,i-1}=-1,\ A_{n-1,n}=-2$.

3 Cayley-Klein structure on quantum symplectic groups $Fun(Sp_q(n;j))$

The Cayley-Klein structure is introduced to quantum symplectic groups $Fun(Sp_q(n))$, as well as in the classical case with the help of the map

$$\psi\hat{x}_k=x_kJ_k,\quad \psi\hat{x}_{k'}=x_{k'}J_k,\quad k=1,\ldots,n, \tag{6}$$

where $\hat{x}_i$ forms the basis of quantum N-dimension symplectic Cayley-Klein space $Sp_q^N(\mathrm{C};\mathrm{j})$ with commutation relations

$$\hat{R}_q(\hat{x}\otimes\hat{x}) = q(\hat{x}\otimes\hat{x}). \tag{7}$$

Parameter j is introduced in matrix T similarly to the classical case. Commutation and additional relations for generators t_{ik} are defined by

$$RT_1(j)T_2(j) = T_2(j)T_1(j)R, \tag{8}$$

and

$$T(j)CT^t(j)C^{-1} = CT^t(j)C^{-1}T(j) = I, \tag{9}$$

whith $T_1(j) = T(j)\otimes I, \quad T_2(j) = I\otimes T(j), \quad C = C_0q^{\rho}$, $\rho = \mathrm{diag}(n, n-1, \ldots, 1, -1, \ldots, -n)$. The coproduct, counit and antipode is defined generally as

$$\triangle(T(j)) = T(j)\dot{\otimes}T(j), \quad \epsilon(T(j)) = I,$$
$$S(T(j)) = CT^t(j)C^{-1}. \tag{10}$$

The "action" δ': $Sp_q^N(\mathrm{C};\mathrm{j}) \to Fun(Sp_q(n;j))\otimes Sp_q^N(\mathrm{C};\mathrm{j})$ of quantum group
$Fun(Sp_q(n;j))$ on quantum symplectic Cayley-Klein space $Sp_q^N(\mathrm{C};j)$, is defined by $\delta' = T(j)\dot{\otimes}\hat{x}$, preserve the bilinear form $\hat{x}^t\dot{\otimes}C\hat{x}$.

4 Quantum algebra $sp_q(n;j)$ as dual to $Fun(Sp_q(n;j))$

Dual to $Fun(Sp_q(n;j))$ the quantum algebra $sp_q(n;j)$ is defined by the relation [2]

$$< L^{(\pm)}(j), T(j) >= R^{(\pm)}.$$
$$R^{(-)} = R_q^{-1} = R_{q^{-1}}, \quad R^{(+)} = PR_qP, \quad Pu\otimes v = v\otimes u. \tag{11}$$

Generators of the algebra $sp_q(n;j)$ have the following form

$$\left(L^{(+)}(j)\right)_{ij} = \tilde{J}^k_m l^{(+)}_{ij}, \quad for\ \ i\le j\ \ and\ \ \left(L^{(+)}(j)\right)_{ij} = 0 \quad for\ \ i>j,$$

$$\left(L^{(-)}(j)\right)_{ij} = \tilde{J}^k_m l^{(-)}_{ij}, \quad for\ \ i\ge j\ \ and\ \ \left(L^{(-)}(j)\right)_{ij} = 0 \quad for\ \ i<j,$$

$$k=i,\ \ i\le n; \quad k=i',\ \ i>n; \quad m=j,\ \ j\le n; \quad m=j',\ \ j>n,$$

$$\tilde{J}^k_m = \prod_{n=min(k,m)}^{max(k,m)-1} j_n^{-1}.$$

Commutation and additional relations for generators of $sp_q(n;j)$ algebras are given by

$$\begin{aligned} R^{(+)}L_1^{(\varepsilon)}(j)L_2^{(\varepsilon)}(j) &= L_2^{(\varepsilon)}(j)L_1^{(\varepsilon)}(j)R^{(+)}, & (12)\\ R^{(+)}L_1^{(+)}(j)L_2^{(-)}(j) &= L_2^{(-)}(j)L_1^{(+)}(j)R^{(+)}, & (13)\\ L(j)C^tL^t(j)(C^{-1})^t &= C^tL^t(j)(C^{-1})^tL(j) = I, & (14)\end{aligned}$$

where $L_1^{(\varepsilon)}(j) = L^{(\varepsilon)}(j) \otimes I, \quad L_2^{(\varepsilon)}(j) = I \otimes L^{(\varepsilon)}(j), \quad \varepsilon = \pm$. Coproduct and counit are defined generally as

$$\triangle(L^{\pm}(j)) = L^{\pm}(j) \dot{\otimes} L^{\pm}(j), \quad \epsilon(L^{\pm}(j)) = I. \tag{15}$$

The isomorphism between algebra $sp_q(n;j)$ and universal enveloping algebra $U_z(sp(n;j))$ is defined by the following expression:

$$l_{ii}^{(+)} = e^{z\tilde{H}_i}, \quad l_{nn}^{(+)} = e^{z\tilde{H}_n}, \tag{16}$$

$$l_{i,i+1}^{(+)} = \lambda q^{-\frac{1}{2}} X_i^+ e^{\frac{z}{2}(\tilde{H}_i + \tilde{H}_{i+1})}, \quad l_{i+1,i}^{(-)} = -\lambda q^{\frac{1}{2}} X_i^- e^{-\frac{z}{2}\widetilde{(\tilde{H}}_i + \tilde{H}_{i+1})} \tag{17}$$

$$l_{n,n+1}^{(+)} = \lambda X_n^+ e^{\frac{z\tilde{H}_n}{2}}, \quad l_{n+1,n}^{(-)} = \lambda X_n^- e^{-\frac{z\tilde{H}_n}{2}}, \tag{18}$$

$$H_i = \tilde{H}_i - \tilde{H}_{i+1}, \quad H_n = 2\tilde{H}_n.$$

Commutation relations in $U_z(sp(n;j))$ are given by

$$[H_i, H_j] = 0, \quad [H_i, X_j^{\pm}] = \pm(\alpha_i, \alpha_j) X_j^{\pm},$$

$$\left[X_i^+, X_j^-\right] = j_i^2 [H_i]_q \delta_{ij} \qquad [H_i]_q = \tfrac{sh(zH_i)}{sh(z)}. \tag{19}$$

Equations (19) defines the structure of universal enveloping algebra $U_z(sp(n;j))$. In the case $j_i = \iota_i$ we have obtained a new nonsemisimple algebra.

5 Case $n = 2$ as example

The quantum symplectic group $Fun(Sp_q(2;j))$ is generated by the matrix with noncommutative generators

$$T(j) = \begin{pmatrix} t_{11} & jt_{12} & jt_{13} & t_{14} \\ jt_{21} & t_{22} & t_{23} & jt_{24} \\ jt_{31} & t_{32} & t_{33} & jt_{34} \\ t_{41} & jt_{42} & jt_{43} & t_{44} \end{pmatrix}. \tag{20}$$

For example in explicit form the antipode is as follows

$$S(T(j)) = \begin{pmatrix} t_{44} & jq^{-1}t_{34} & -jq^{-3}t_{24} & -q^{-4}t_{14} \\ jqt_{43} & t_{33} & -q^{-2}t_{23} & -jq^{-3}t_{13} \\ -jq^3t_{42} & -q^2t_{32} & t_{22} & jq^{-1}t_{12} \\ -q^4t_{41} & -jq^3t_{31} & jqt_{21} & t_{11} \end{pmatrix}. \tag{21}$$

The generators of the dual to $Fun(Sp_q(2;j))$ quantum algebra $sp_q(2;j)$ written in compact matrix form as

$$L^{(+)}(j) = \begin{pmatrix} l_{11}^{(+)} & j^{-1}l_{12}^{(+)} & j^{-1}l_{13}^{(+)} & l_{14}^{(+)} \\ 0 & l_{22}^{(+)} & l_{23}^{(+)} & j^{-1}l_{24}^{(+)} \\ 0 & 0 & l_{33}^{(+)} & j^{-1}l_{34}^{(+)} \\ 0 & 0 & 0 & l_{44}^{(+)} \end{pmatrix}, \tag{22}$$

$$L^{(-)}(j) = \begin{pmatrix} l_{11}^{(-)} & 0 & 0 & 0 \\ j^{-1}l_{21}^{(-)} & l_{22}^{(-)} & 0 & 0 \\ j^{-1}l_{31}^{(-)} & l_{32}^{(-)} & l_{33}^{(-)} & 0 \\ l_{41}^{(-)} & j^{-1}l_{42}^{(-)} & j^{-1}l_{43}^{(-)} & l_{44}^{(-)} \end{pmatrix}. \tag{23}$$

Commutation and additional relations for generators of $sp_q(2;j)$ algebras are given by

$$\begin{gathered}
ql_{11}^{(+)}l_{12}^{(+)} = l_{12}^{(+)}l_{11}^{(+)}, \quad l_{11}^{(+)}l_{23}^{(+)} = l_{23}^{(+)}l_{11}^{(+)}, \quad l_{11}^{(+)}l_{32}^{(-)} = l_{32}^{(-)}l_{11}^{(+)}, \\
l_{11}^{(+)}l_{21}^{(-)} = ql_{21}^{(-)}l_{11}^{(+)}, \quad l_{11}^{(\pm)}l_{kk}^{(\pm)} = l_{kk}^{(\pm)}l_{11}^{(\pm)}, \quad k = 1, \ldots, 4, \\
ql_{22}^{(+)}l_{21}^{(-)} = l_{21}^{(-)}l_{22}^{(+)}, \quad l_{22}^{(+)}l_{32}^{(-)} = q^2 l_{32}^{(-)}l_{22}^{(+)}, \quad l_{22}^{(+)}l_{12}^{(+)} = ql_{12}^{(+)}l_{22}^{(+)}, \\
q^2 l_{22}^{(+)}l_{23}^{(+)} = l_{23}^{(+)}l_{22}^{(+)}, \quad ql_{23}^{(+)}l_{12}^{(+)} - l_{12}^{(+)}l_{23}^{(+)} = \tilde{\lambda} l_{13}^{(+)}l_{22}^{(+)}, \quad \tilde{\lambda} = q^2 - q^{-2}, \\
[l_{12}^{(+)}, l_{21}^{(-)}] = \lambda(l_{11}^{(+)}l_{22}^{(-)} - l_{11}^{(-)}l_{22}^{(+)}), \quad [l_{23}^{(+)}, l_{32}^{(-)}] = \tilde{\lambda}(l_{22}^{(+)}l_{22}^{(+)} - l_{22}^{(-)}l_{22}^{(-)}), \\
ql_{32}^{(-)}l_{21}^{(-)} - l_{21}^{(-)}l_{32}^{(-)} = \tilde{\lambda} l_{31}^{(-)}l_{22}^{(-)}, \quad l_{12}^{(+)}l_{32}^{(-)} = ql_{32}^{(-)}l_{12}^{(+)}, \quad ql_{23}^{(+)}l_{21}^{(-)} = l_{21}^{(-)}l_{23}^{(+)}.
\end{gathered} \tag{24}$$

Antipode :

$$S(L^{(+)}(j)) = \begin{pmatrix} l_{44}^{(+)} & j^{-1}ql_{34}^{(+)} & -j^{-1}q^3 l_{24}^{(+)} & -q^4 l_{14}^{(+)} \\ 0 & l_{33}^{(+)} & -q^2 l_{23}^{(+)} & -j^{-1}q^3 l_{13}^{(+)} \\ 0 & 0 & l_{22}^{(+)} & j^{-1}ql_{12}^{(+)} \\ 0 & 0 & 0 & l_{11}^{(+)} \end{pmatrix}, \tag{25}$$

$$S(L^{(-)}(j)) = \begin{pmatrix} l_{44}^{(-)} & 0 & 0 & 0 \\ j^{-1}q^{-1}l_{43}^{(-)} & l_{33}^{(-)} & 0 & 0 \\ -j^{-1}q^{-3}l_{42}^{(-)} & -q^{-2}l_{32}^{(-)} & l_{22}^{(-)} & 0 \\ -q^{-4}l_{41}^{(-)} & -j^{-1}q^{-3}l_{31}^{(-)} & j^{-1}q^{-1}l_{21}^{(-)} & l_{11}^{(-)} \end{pmatrix}. \tag{26}$$

The isomorphism between quantum algebra $sp_q(2;j)$ and universal enveloping algebra $U_z(sp(2;j))$ is defined by the following expressions ($q = e^{\varepsilon z}, \ \varepsilon = \pm 1$):

$$l_{11}^{(+)} = e^{z\tilde{H}_i}, \quad l_{22}^{(+)} = e^{z\tilde{H}_n}, \tag{27}$$

$$l_{12}^{(+)} = \lambda q^{-\frac{1}{2}} X_1^+ e^{\frac{z}{2}(\tilde{H}_1 + \tilde{H}_2)}, \quad l_{21}^{(-)} = -\lambda q^{\frac{1}{2}} X_1^- e^{-\frac{z}{2}(\widetilde{H_i} + \tilde{H}_2)}, \tag{28}$$

$$l_{23}^{(+)} = \lambda q^{-\frac{1}{2}} X_2^+ e^{\frac{z\tilde{H}_2}{2}}, \quad l_{32}^{(-)} = \lambda q^{\frac{1}{2}} X_2^- e^{-\frac{z\tilde{H}_2}{2}}, \tag{29}$$

$$H_1 = \tilde{H}_1 - \tilde{H}_2, \quad H_2 = 2\tilde{H}_2.$$

Other generators may be obtained from additional relation (14)

$$\begin{aligned}
l_{13}^{(+)} &= (\tilde{\lambda})^{-1}(ql_{23}^{+}l_{12}^{(+)} - l_{12}^{(+)}l_{23}^{(+)})l_{22}^{(-)}, \quad l_{31}^{(-)} = (\tilde{\lambda})^{-1}(ql_{32}^{(-)}l_{21}^{(-)} - l_{21}^{(-)}l_{32}^{(-)})l_{22}^{(+)}, \\
l_{14}^{(+)} &= j^{-2}\lambda^{-1}(l_{13}^{(+)}l_{12}^{(+)} - q^2 l_{12}^{(+)}l_{13}^{(+)})l_{11}^{(-)}, \qquad l_{34}^{+} = -l_{12}^{(+)}l_{11}^{(-)}l_{22}^{(-)}, \\
l_{41}^{(-)} &= j^{-2}\lambda^{-1}(l_{31}^{(-)}l_{21}^{(-)} - q^2 l_{21}^{(+)}l_{31}^{(+)})l_{11}^{(+)}, \qquad l_{43}^{-} = -l_{21}^{(-)}l_{11}^{(+)}l_{22}^{(+)}, \\
l_{ii}^{(+)} &= l_{i'i'}^{(-)}, \qquad l_{ii}^{(-)} = (l_{ii}^{(+)})^{-1}, \qquad i = 1, \ldots, 4.
\end{aligned}$$

Commutation relations in $U_z(sp(2;j))$ are given by

$$\begin{aligned}
[H_i, H_j] = 0, \quad [H_i, X_j^{\pm}] &= \pm(\alpha_i, \alpha_j)X_j^{\pm}, \\
\left[X_i^+, X_j^-\right] &= j_i^2 [H_i]_q \delta_{ij}.
\end{aligned} \tag{30}$$

$$(\alpha_i, \alpha_j) = -2\varepsilon \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix}. \tag{31}$$

In the case $j = \iota$ we have obtained a new nonsemisimple algebra.

References

[1] N.A. Gromov, *Contractions and analytical continuations of classical groups. Unified approach*, Syktyvkar, mi SC, 1990 (in Russian).

[2] N.Yu. Reshetikhin, L.A. Takhtajan, L.D. Faddeev, *Algebra and analysis.* (1989), Vol. 1, No.1, pp. 178-206.